

3. (Amended) The cell according to claim 1, wherein said ligand-responsive transcription control factor is one selected from an aryl hydrocarbon receptor, intranuclear hormone receptor, estrogen receptor, androgen receptor and thyroid hormone receptor.

4. (Amended) The cell according to claim 1, wherein said ligand-responsive transcription control factor is an aryl hydrocarbon receptor.

9. (Amended) An animal cell expressing an aryl hydrocarbon receptor and an Arnt receptor and securely maintaining a DNA comprising in a molecule, the following genes (a) and (b):

(a) a reporter gene connected downstream from a transcription control region, wherein said transcription control region substantially consists of a recognition sequence of said aryl hydrocarbon receptor and a minimum promoter which can function in said cell and

(b) a selective marker gene which can function in said cell;

provided that the following gene (c):

(c) a reporter gene connected downstream from a promoter which transcription activity is unchanged by having a

ligand-responsive transcription control factor contacted with a ligand of said ligand-responsive transcription control factor, said reporter gene (c) coding a protein which can be differentiated from the protein coded by said gene (a)

is not present in said cell.

sub 24 14. (Amended) A method for obtaining an animal cell for measuring the ability to control the activity of a ligand-responsive transcription control factor, said method comprising:
(i) introducing into an animal cell, a DNA comprising in a molecule the following genes (a) and (b):

- a3*
- (a) a reporter gene connected downstream from a transcription control region, wherein said transcription control region substantially consists of a recognition sequence of said ligand-responsive transcription control factor and a minimum promoter which can function in said cell, and
 - (b) a selective marker gene which can function in said cell,

said animal cell being

an animal cell that comprises a DNA comprising a gene coding the ligand-responsive control factor introduced thereto before, after or during the same time of above

step (i) or that naturally has an ability to express the gene coding the ligand-responsive transcription control factor,

provided that a reporter gene (c) connected downstream from a promoter which transcription activity is unchanged by having said responsive transcription control factor contacted with a ligand of said ligand-responsive transcription control factor, said reporter gene (c) coding a protein which can be differentiated from the protein coded by said gene (a), is not present in the cell; and

(ii) recovering from the transformed cell obtained from step (i), a transformed cell having said introduced DNA securely maintained therein.
